

# Literaturverzeichnisse 2017

## **Inhaltsverzeichnis**

[04-2017: Literaturverzeichnis Sonntag](#)

[04-2017: Literaturverzeichnis Jerosch](#)

[04-2017: Literaturverzeichnis Günther](#)

[04-2017: Literaturverzeichnis Dynbil](#)

[05-2017: Literaturverzeichnis von Rüden](#)

[06-2017: Literaturverzeichnis Hempfling](#)

[06-2017: Literaturverzeichnis Keshmiri](#)

[06-2017: Literaturverzeichnis Wagner](#)

# Vollständige Literatur zu Robert Sonntag et al.: Die Qual der Wahl: Welche Hüft-Gleitpaarung für die orthopädische Praxis? OUP 04-2017

## Literatur

- Affatato S, Leardini W, Jedenmalm A, Ruggeri O, Toni A: Larger diameter bearings reduce wear in metal-on-metal hip implants. *Clin Orthop Relat Res* 2007; 456: 153–158
- Affatato S, Spinelli M, Zavalloni M, Traina F, Carmignato S, Toni A: Ceramic-On-Metal for Total Hip Replacement: Mixing and Matching Can Lead to High Wear. *Artificial Organs* 2010; 34: 319–323
- Australian National Joint Replacement Registry. Annual Report. 2013
- Australian National Joint Replacement Registry. Annual Report. 2016
- Bergmann G, Bender A, Dymke J, Duda G, Damm P: Standardized Loads Acting in Hip Implants. *PLoS One* 2016; 11: e0155612
- Bragdon CR, Doerner M, Martell J, Jarrett B, Palm H, Malchau H: The 2012 John Charnley Award: Clinical multicenter studies of the wear performance of highly crosslinked remelted polyethylene in THA. *Clin Orthop Relat Res* 2013; 471: 393–402
- Clarke IC, Gustafson A, Jung H, Fujisawa A: Hip-simulator ranking of polyethylene wear: comparisons between ceramic heads of different sizes. *Acta Orthop Scand* 1996; 67: 128–132
- Cole JC, Lemons JE, Eberhardt AW: Gamma irradiation alters fatigue-crack behavior and fracture toughness in 1900H and GUR 1050 UHMWPE. *J Biomed Mater Res* 2002; 63: 559–566
- Damm P, Graichen F, Rohlmann A, Bender A, Bergmann G: Total hip joint prosthesis for in vivo measurement of forces and moments. *Med Eng Phys* 2010; 32: 95–100
- Dobbs HS, Minski MJ: Metal ion release after total hip replacement. *Biomaterials* 1980; 1: 193–198
- Dowson D, Hardaker C, Flett M, Isaac GH: A hip joint simulator study of the performance of metal-on-metal joints: Part II: Design. *J Arthroplasty* 2004; 19 (8, Supplement 1): 124–130
- EFORT, Arbeitsgemeinschaft Endoprothetik, Deutsche Arthroshilfe e.V.: Aktuelle Konsens-Empfehlungen zur Handhabung von Metall-Metall-Gleitpaarungen. 2012
- Endo M, Tipper JL, Barton DC, Stone MH, Ingham E, Fisher J: Comparison of wear, wear debris and functional biological activity of moderately crosslinked and non-crosslinked polyethylenes in hip prostheses. *Proc Inst Mech Eng H* 2002; 216: 111–122
- Epinette JA, Manley MT: No differences found in bearing related hip survivorship at 10–12 years follow-up between patients with ceramic on highly crosslinked polyethylene bearings compared to patients with ceramic on ceramic bearings. *J Arthroplasty* 2014; 29: 1369–1372
- Fisher J: Bioengineering reasons for the failure of metal-on-metal hip prostheses: an engineer's perspective. *J Bone Joint Surg Br* 2011; 93-B: 1001–1004
- Galvin A, Brockett C, Williams S et al.: Comparison of wear of ultra-high molecular weight polyethylene acetabular cups against surface-engineered femoral heads. *Proc Inst Mech Eng H* 2008; 222: 1073–1080
- Galvin AL, Jennings LM, Tipper JL, Ingham E, Fisher J: Wear and creep of highly crosslinked polyethylene against cobalt chrome and ceramic femoral heads. *Proc Inst Mech Eng H* 2010; 224: 1175–1183
- Geerdink CH, Grimm B, Ramakrishnan R, Rondhuis J, Verburg AJ, Tonino AJ: Crosslinked polyethylene compared to conventional polyethylene in total hip replacement: pre-clinical evaluation, in-vitro testing and prospective clinical follow-up study. *Acta Orthop* 2006; 77: 719–725
- Gutmanas EY, Gotman I: PIRAC Ti nitride coated Ti-6Al-4V head against UHMWPE acetabular cup-hip wear simulator study. *J Mater Sci Mater Med* 2004; 15: 327–330
- Habermann B, Ewald W, Rauschmann M, Zichner L, Kurth AA: Fracture of ceramic heads in total hip replacement. *Arch Orthop Trauma Surg* 2006; 126: 464–470
- Hamilton WG, McAuley JP, Dennis DA, Murphy JA, Blumenfeld TJ, Politi J: THA With Delta Ceramic on Ceramic: Results of a Multicenter Investigational Device Exemption Trial. *Clin Orthop Relat Res* 2010; 468: 358–366
- Harman MK, Banks SA, Hodge WA: Wear analysis of a retrieved hip implant with titanium nitride coating. *J Arthroplasty* 1997; 12: 938–945
- Hartmann A, Hannemann F, Lutzner J et al.: Metal ion concentrations in body fluids after implantation of hip replacements with metal-on-metal bearing – systematic review of clinical and epidemiological studies. *PLoS One* 2013; 8: e70359
- Hatton A, Nevelos JE, Nevelos AA, Banks RE, Fisher J, Ingham E: Alumina-alumina artificial hip joints. Part I: a histological analysis and characterisation of wear debris by laser capture microdissection of tissues retrieved at revision. *Biomaterials* 2002; 23: 3429–3440
- Hermida JC, Bergula A, Chen P, Colwell CW, Jr, D'Lima DD: Comparison of the wear rates of twenty-eight and thirty-two-millimeter femoral heads on cross-linked polyethylene acetabular cups in a wear simulator. *J Bone Joint Surg Am* 2003; 85-a: 2325–2331
- Howie DW, Holubowycz OT, Middleton R: Large femoral heads decrease the incidence of dislocation after total hip arthroplasty: a randomized controlled trial. *J Bone Joint Surg Am* 2012; 94: 1095–1102
- Ingram JH, Stone M, Fisher J, Ingham E: The influence of molecular weight, crosslinking and counterface roughness on TNF-alpha production by macrophages in response to ultra high molecular weight polyethylene particles. *Biomaterials* 2004; 25: 3511–3522
- Kadar T, Hallan G, Aamodt A et al.: Wear and migration of highly cross-linked and conventional cemented polyethylene cups with cobalt chrome or Oxinium femoral heads: a randomized radiostereometric study of 150 patients. *J Orthop Res* 2011; 29: 1222–1229
- Koo KH, Ha YC, Jung WH, Kim SR, Yoo JJ, Kim HJ: Isolated fracture of the ceramic head after third-generation alumina-on-alumina total hip arthroplasty. *J Bone Joint Surg Am* 2008; 90: 329–336
- Langton DJ, Jameson SS, Joyce TJ, Hallab NJ, Natsu S, Nargol AVF: Early failure of metal-on-metal bearings in hip resurfacing and large-diameter total hip replacement: A consequence of excess wear. *J Bone Joint Surg Br* 2010; 92-B: 38–46
- Losina E, Barrett J, Mahomed NN, Baron JA, Katz JN: Early failures of total hip replacement: effect of surgeon volume. *Arthritis Rheum* 2004; 50: 1338–1343

32. Mai K, Verioti C, Ezzet K, Copp S, Walker R, Colwell C: Incidence of 'Squeaking' After Ceramic-on-Ceramic Total Hip Arthroplasty. *Clin Orthop Relat Res* 2009; 468: 413–417
33. Manley M, Ong K, Lau E, Kurtz SM: Effect of volume on total hip arthroplasty revision rates in the United States Medicare population. *J Bone Joint Surg Am* 2008; 90: 2446–2451
34. Massin P, Lopes R, Masson B, Mainard D: Does Biolox Delta ceramic reduce the rate of component fractures in total hip replacement? *Orthop Traumatol Surg Res* 2014; 100 (6 Suppl): 317–321
35. Morison ZA, Patil S, Khan HA, Bogoch ER, Schemitsch EH, Waddell JP: A randomized controlled trial comparing Oxinium and cobalt-chrome on standard and cross-linked polyethylene. *J Arthroplasty* 2014; 29 (9 Suppl): 164–168
36. Mutimer J, Devane PA, Adams K, Horne JG: Highly crosslinked polyethylene reduces wear in total hip arthroplasty at 5 years. *Clin Orthop Relat Res* 2010; 468: 3228–3233
37. National Joint Registry for England Wales, Northern Ireland and the Isle of Man. Annual Report. 2015
38. Owen DH, Russell NC, Smith PN, Walter WL: An estimation of the incidence of squeaking and revision surgery for squeaking in ceramic-on-ceramic total hip replacement: a meta-analysis and report from the Australian Orthopaedic Association National Joint Registry. *Bone Joint J* 2014; 96-B: 181–187
39. Pandit H, Glyn-Jones S, McLardy-Smith P et al.: Pseudotumours associated with metal-on-metal hip resurfacings. *J Bone Joint Surg Br* 2008; 90: 847–851
40. Pappas MJ, Makris G, Buechel FF: Titanium Nitride Ceramic Film Against Polyethylene: A 48 Million Cycle Wear Test. *Clin Orthop Relat Res*. 1995; 317: 64–70
41. Park YS, Hwang SK, Choy WS, Kim YS, Moon YW: Ceramic Failure After Total Hip Arthroplasty with an Alumina-on-Alumina Bearing. *J Bone Joint Surg Am* 2006; 88: 780–787
42. Prokopetz JJ, Losina E, Bliss RL, Wright J, Baron JA, Katz JN: Risk factors for revision of primary total hip arthroplasty: a systematic review. *BMC Musculoskelet Disord* 2012; 13: 251
43. Purdue PE, Koulouvaris P, Nestor BJ, Sulco TP: The central role of wear debris in periprosthetic osteolysis. *HSS J* 2006; 2: 102–113
44. Raimondi MT, Pietrabissa R: The in-vivo wear performance of prosthetic femoral heads with titanium nitride coating. *Biomaterials* 2000; 21: 907–913
45. Saikko V, Ahlroos T, Revitzer H, Rytö O, Kuosmanen P: The effect of acetabular cup position on wear of a large-diameter metal-on-metal prosthesis studied with a hip joint simulator. *Tribol Int* 2013; 60: 70–76
46. Savarino L, Baldini N, Ciapetti G, Pellacani A, Giunti A: Is wear debris responsible for failure in alumina-on-alumina implants? *Acta Orthop* 2009; 80: 162–167
47. Smith AJ, Dieppe P, Vernon K, Porter M, Blom AW: Failure rates of stemmed metal-on-metal hip replacements: analysis of data from the National Joint Registry of England and Wales. *Lancet* 2012; 379: 1199–1204
48. Sonntag R, Reinders J, Müller U, Kretzer JP: Wahl der richtigen Gleitpaarung in der Hüftendoprothetik. *Z Orthop Unfall* 2015; 153: 587–596
49. Tower SS, Currier JH, Currier BH, Lyford KA, Van Citters DW, Mayor MB: Rim cracking of the cross-linked longevity polyethylene acetabular liner after total hip arthroplasty. *J Bone Joint Surg Am* 2007; 89: 2212–2217
50. Traina F, Tassinari E, De Fine M, Bordini B, Toni A: Revision of Ceramic Hip Replacements for Fracture of a Ceramic Component. *AAOS Exhibit Selection* 2011; 93: e147
51. Traina F, De Fine M, Di Martino A, Faldini C: Fracture of Ceramic Bearing Surfaces following Total Hip Replacement: A Systematic Review. *Biomed Res Int* 2013; 2013: 157247
52. Waewsawangwong W, Goodman SB: Unexpected Failure of Highly Cross-Linked Polyethylene Acetabular Liner. *J Arthroplasty* 2012; 27: 323.e1–323.e4
53. Weiss C, Gdaniec P, Hoffmann NP, Hothan A, Huber G, Morlock MM: Squeak in hip endoprosthesis systems: An experimental study and a numerical technique to analyze design variants. *Med Eng Phys* 2010; 32: 604–609
54. Willert HG, Buchhorn GH, Fayyazi A et al.: Metal-on-Metal Bearings and Hypersensitivity in Patients with Artificial Hip Joints A Clinical and Histomorphological Study. *J Bone Joint Surg Am* 2005; 87: 28–36
55. Zietz C, Fabry C, Middelborg L, Fulda G, Mittelmeier W, Bader R: Wear testing and particle characterisation of sequentially crosslinked polyethylene acetabular liners using different femoral head sizes. *J Mater Sci Mater Med* 2013; 24: 2057–2065
56. Zietz C, Fabry C, Reinders J et al.: Wear testing of total hip replacements under severe conditions. *Expert Rev Med Devices* 2015; 12: 393–410

# Vollständige Literatur zu Jörg Jerosch et al.: Welche Ergebnisse sind mit einem endoskopischen Release der Plantarfaszie (ERPF) zu erreichen? OUP 04-2017

## Literatur

1. Acevedo JI, Beskin JL: Complications of plantar fascia rupture associated with corticosteroid injection. *Foot Ankle Int* 1998; 19: 91–7
2. Arangio GA, Chen C, Kim W: Effect of cutting the plantar fascia on mechanical properties of the foot. *Clin Orthop* 1997; 339: 227–31
3. Barrett SL, Day SV, Pignetti TT, Robinson LB: Endoscopic plantar fasciotomy: a multi-surgeon prospective analysis of 652 cases. *J Foot Ankle Surg* 1995; 34: 400–6
4. Barrett SL, Day SV: Endoscopic plantar fasciotomy for chronic plantar fasciitis/heel spur syndrome: surgical technique – early clinical results. *J Foot Surg* 1991; 30: 568–70
5. Basford JR, Malanga GA, Krause DA, Harmsen WS: A randomized controlled evaluation of low-intensity laser therapy: plantar fasciitis. *Arch Phys Med Rehabil* 1998; 79: 249–54
6. Batt ME, Tanji JL, Skattum N: Plantar fasciitis: a prospective randomized clinical trial of the tension night splint. *Clin J Sport Med* 1996; 6: 158–62
7. Bedi HS, Love BR: Differences in impulse distribution in patients with plantar fasciitis. *Foot Ankle Int* 1998; 19: 153–156
8. Brekke MK, Green DR: Retrospective analysis of minimal-incision, endoscopic, and open procedures for heel spur syndrome. *J Am Podiatr Med Assoc* 1998; 88: 64–72
9. Brugh AM, Fallat LM, Savoy-Moore RT: Lateral column symptomatology following plantar fascial release: a prospective study. *J Foot Ankle Surg* 2002; 41: 365–71
10. Cardinal E, Chhem RK, Beaugard CG, Aubin B, Pelletier M: Plantar fasciitis: sonographic evaluation. *Radiology* 1996; 201: 257–59
11. Chandler TJ: Iontophoresis of 0.4 % dexamethasone for plantar fasciitis. *Clin J Sport Med* 1998; 8: 68
12. Chandler TJ, Kibler WB: A biomechanical approach to the prevention, treatment and rehabilitation of plantar fasciitis. *Sports Med* 1993; 15: 344–52
13. Daly PJ, Kitaoka HB, Chao EY: Plantar fasciotomy for intractable plantar fasciitis: clinical results and biomechanical evaluation. *Foot Ankle* 1992; 13: 188–95
14. de Souza H, Reed L: An inexpensive „orthosis“ for plantar fasciitis. *Med J Aust* 1997; 167: 509
15. Gill LH, Kiezbak GM: Outcome of non-surgical treatment for plantar fasciitis. *Foot Ankle Int* 1996; 17: 527–32
16. Graves RH 3rd, Levin DR, Giacomelli J, White PR, Russell RD: Fluoroscopy-assisted plantar fasciotomy and calcaneal exostectomy: a retrospective study and comparison of surgical techniques. *J Foot Ankle Surg* 1994; 33: 475–81
17. Hawkins BJ, Langermen RJ Jr, Gibbons T, Calhoun JH: An anatomic analysis of endoscopic plantar fascia release. *Foot Ankle Int* 1995; 16: 552–558
18. Heller KD: Extracorporeal shockwave therapy in heel spur – analysis of the literature. *Z Orthop Ihre Grenzgeb* 1999; 137: 13–5
19. Jerosch J: Endoscopic release of plantar fasciitis – a benign procedure? *Foot Ankle Int* 2000; 21: 511–13
20. Jerosch J, Schunck J, Liebsch D, Filler T: Indication, surgical technique and results of endoscopic fascial release in plantar fasciitis (EFRPF). *Knee Surg Sports Traumatol. Arthrosc.* 2004; 12: 471–77
21. Kane D, Greaney T, Bresnihan B, Gibney R, FitzGerald O: Ultrasound guided injection of recalcitrant plantar fasciitis. *Ann Rheum Dis* 1998; 57: 383–84
22. Kibler WB, Goldberg C, Chandler TJ: Functional biomechanical deficits in running athletes with plantar fasciitis. *Am J Sports Med* 1991; 19: 66–71
23. Kier R: Magnetic resonance imaging of plantar fasciitis and other causes of heel pain. *Magn Reson Imaging Clin N Am* 1994; 2: 97–107
24. Kinley S, Frascone S, Calderone D, Wertheimer SJ, Squire MA, Wiseman FA: Endoscopic plantar fasciotomy versus traditional heel spur surgery: a prospective study. *J Foot Ankle Surg* 1993; 32: 595–603
25. Krischek O, Rompe JD, Herbsthofner B, Nafe B: Symptomatic low-energy shockwave therapy in heel pain and radiologically detected plantar heel spur. *Z Orthop* 1998; 136: 169–74
26. Maier M, Durr HR, Kohler S, Staupendahl D, Pfahler M, Refior HJ, Meier M: Analgesic effect of low energy extracorporeal shock waves in tendinosis calcarea, epicondylitis humeri radialis and plantar fasciitis. *Z Orthop Ihre Grenzgeb* 2000; 138: 34–8
27. Manoli A 2nd, Harper MC, Fitzgibbons TC, McKernan DJ: Calcaneal fracture after cortical bone removal. *Foot Ankle* 1992; 13: 523–25
28. Mizel MS, Marymont JV, Trepman E: Treatment of plantar fasciitis with a night splint and shoe modification consisting of a steel shank and anterior rocker bottom. *Foot Ankle Int* 1996; 17: 732–35
29. Murphy GA, Pneumaticos SG, Kamaric E, Noble PC, Trevino SG, Baxter DE: Biomechanical consequences of sequential plantar fascia release. *Foot Ankle Int* 1998; 19: 149–52
30. Ogilvie-Harris DJ, Mahomed N, Demazière A: Anterior impingement of the ankle treated by arthroscopic removal of bony spurs. *J Bone Joint Surg Br* 1993; 75: 437–40
31. Placzek R, Deuretzbacher G, Meiss AL: Treatment of chronic plantar fasciitis with Botulinum toxin A: preliminary clinical results. *Clin J Pain* 2006; 22: 190–2
32. Placzek R, Hölscher A, Deuretzbacher G, Meiss L, Perka C: Treatment of chronic plantar fasciitis with botulinum toxin A – an open pilot study on 25 patients with a 14-week-follow-up. *Z Orthop Ihre Grenzgeb* 2006; 144: 405–9
33. Powell M, Post WR, Keener J, Wearden S: Effective treatment of chronic plantar fasciitis with dorsiflexion night splints: a crossover prospective randomized outcome study. *Foot Ankle Int* 1998; 19: 10–8
34. Reeve F, Laughlin RT, Wright DG: Endoscopic plantar fascia release: a cross-sectional anatomic study. *Foot Ankle Int* 1997; 18: 398–401
35. Rompe JD, Cacchio A, Weil L Jr et al.: Plantar fascia-specific stretching versus radial shock-wave therapy as initial treatment of plantar fasciopathy. *J Bone Joint Surg Am* 2010; 92: 2514–22
36. Sadat-Ali M: Plantar fasciitis/calcaneal spur among security forces personnel. *Mil Med* 1998; 163: 56–7
37. Sammarco GJ, Helfrey RB: Surgical treatment of recalcitrant plantar fasciitis. *Foot Ankle Int* 1996; 17: 520–526
38. Sammarco GJ, Idusuyi OB: Stress fracture of the base of the third metatarsal after an endoscopic plantar fasciotomy: a

- case report. *Foot Ankle Int* 1998; 19: 157–159
39. Schon LC, Baxter DE: Neuropathies of the foot and ankle in athletes. *Clin Sports Med*. 1990; 9: 489–509.
40. Sellman JR: Plantar fascia rupture associated with corticosteroid injection. *Foot Ankle Int* 1994; 15: 376–81
41. Smith WK, Noriega JA, Smith WK Jr: Resection of a plantar calcaneal spur using the holmium:yttrium-aluminum-garnet (Ho:YAG) laser. *J Am Podiatr Med Assoc* 2001; 91: 142–6
42. Sollitto RJ, Plotkin EL, Klein PG, Mullin P: Early clinical results of the use of radiofrequency lesioning in the treatment of plantar fasciitis. *J Foot Ankle Surg* 1997; 36: 215–19
43. Thordarson DB, Kumar PJ, Hedman TP, Ebrahimzadeh E: Effect of partial versus complete plantar fasciotomy on the windlass mechanism. *Foot Ankle Int* 1997; 18: 16–20
44. Tomczak RL, Haverstock BD: A retrospective comparison of endoscopic plantar fasciotomy to open plantar fasciotomy with heel spur resection for chronic plantar fasciitis/heel spur syndrome. *J Foot Ankle Surg* 1995; 34: 305–11
45. Tudor GR, Finlay D, Allen MJ, Belton I: The role of bone scintigraphy and plain radiography in intractable plantar fasciitis. *Nucl Med Commun* 1997; 18: 853–6
46. Vohra PK, Giorgini RJ, Sobel E, Japour CJ, Villalba MA, Rostkowski T: Long-term follow-up of heel spur surgery. A 10-year retrospective study. *J Am Podiatr Med Assoc* 1999; 89: 81–8
47. Wall JR, Harkness MA, Crawford A: Ultrasound diagnosis of plantar fasciitis. *Foot Ankle* 1993; 14: 465–70
48. White AD: An inexpensive „orthosis“ for plantar fasciitis. *Med J Aust* 1997; 166: 616
49. Wolgin M, Cook C, Graham C, Mauldin D: Conservative treatment of plantar heel pain: long-term follow-up. *Foot Ankle Int* 1994; 15: 97–102

# Vollständige Literatur zu Klaus-Peter Günther et al.: Wann ist der richtige Zeitpunkt für die Primäroperation in der Hüftendoprothetik? OUP 04-2017

## Literatur

- Balck F, Lippmann M, Jeszenszky C, Günther KP, Kirschner S: The Influence of Optimism on Functionality after Total Hip Replacement Surgery. *J Health Psychol* 2015 Jan 20. pii: 1359105314566256. [Epub ahead of print]
- Bleß H, Kip M (Hrsg.): Weißbuch Gelenkersatz. Versorgungssituation endoprothetischer Hüft- und Knieoperationen in Deutschland. Springer 2016
- Bozic KJ, Lau E, Kurtz S et al.: Patient-related risk factors for periprosthetic joint infection and postoperative mortality following total hip arthroplasty in Medicare patients. *J Bone Joint Surg Am* 2012; 94: 794–800
- Bozic KJ, Lau E, Ong K et al.: Risk factors for early revision after primary total hip arthroplasty in Medicare patients. *Clin Orthop Relat Res* 2014; 472: 449–54
- Clement ND, Muzammil A, Macdonald D, Howie CR, Biant LC: Socioeconomic status affects the early outcome of total hip replacement. *The Journal of bone and joint surgery* 2011; British volume 93: 464–69
- Deutsches Institut für Medizinische Dokumentation und Information (DIMDI) (Hrsg.) Internationale Klassifikation der Funktionsfähigkeit, Behinderung und Gesundheit (ICF), Genf: WHO, 2005
- Dreinhöfer KE, Dieppe P, Stürmer T et al.: Indications for total hip replacement: comparison of assessments of orthopaedic surgeons and referring physician. *Ann Rheum Dis* 2006; 65: 1346–50
- Dowsey MM, Nikpour M, Choong PF: 2014 Outcomes following large joint arthroplasty: does socio-economic status matter? *BMC Musculoskelet Disor* 2014; 15: 148
- Finkenstädt V, Niehaus F: Die Aussagekraft von Länderrankings im Gesundheitsbereich. Köln: Wissenschaftliches Institut der PKV, 2015
- Fogel GR, Esses SI: Hip spine syndrome: management of coexisting radiculopathy and arthritis of the lower extremity. *The Spine Journal* 2003; 3: 238–41
- Gossec L, Paternotte S, Maillefert JF: et al: The role of pain and functional impairment in the decision to recommend total joint replacement in hip and knee osteoarthritis: an international cross-sectional study of 1909 patients. Report of the OARSI-OMERACT Task Force on total joint replacement. *Osteoarthritis Cartilage* 2011; 19: 147–54
- Greene ME, Rolfson O, Nemes S, Gordon M, Malchau H, Garellick G: Education attainment is associated with patient-reported outcomes: findings from the Swedish Hip Arthroplasty Register. *Clin Orthop Relat Res* 2014; 472: 1868–76
- Günther KP, Haase E, Lange T et al.: Persönlichkeitsprofil und Komorbidität: Gibt es den „schwierigen Patienten“ in der primären Hüftendoprothetik? *Orthopäde* 2015; 44: 555–65
- Haanstra TM, van den Berg T, Ostelo RW et al.: Systematic review: do patient expectations influence treatment outcomes in total knee and total hip arthroplasty? *Health Qual Life Outcomes* 2012; 10: 152
- Haase E, Kopkow C, Beyer F et al.: Patient-reported outcomes and outcome predictors after primary total hip arthroplasty: results from the Dresden Hip Surgery Registry. *Hip Int* 2015; Oct 6:0. doi: 10.5301/hipint.5000300. [Epub ahead of print]
- Hamann C, Günther KP, Lütznier J, Beyer F, Kirschner S: Diabetes mellitus – Ein unterschätzter Risikofaktor in der Hüftendoprothetik? *Osteologie* 2015; (article in press)
- Hamilton DF, Lane JV, Gaston P et al.: What determines patient satisfaction with surgery? A prospective cohort study of 4709 patients following total joint replacement. *BMJ Open* 2013; 3(4). pii: e002525. doi: 10.1136/bmjopen-2012-002525. Print 2013
- Haverkamp D, Klinkenbijl MN, Somford MP, Albers GH, van der Vis HM: Obesity in total hip arthroplasty – does it really matter? A meta-analysis. *Acta Orthop* 2011; 82: 417–22
- Hawker GA, Badley EM, Borkhoff CM et al.: Which patients are most likely to benefit from total joint arthroplasty? *Arthritis Rheum* 2013; 65: 1243–52
- Herberts P, Malchau H: Long-term registration has improved the quality of hip replacement: A review of the Swedish THR Register comparing 160,000 cases. *Acta Orthop* 2000; 71: 111–21
- Hooper GJ, Rothwell AG, Hooper NM, Frampton C: The relationship between the American Society Of Anesthesiologists physical rating and outcome following total hip and knee arthroplasty: an analysis of the New Zealand Joint Registry. *J Bone Joint Surg Am* 2012; 94: 1065–70
- Hutchings A, Neuburger J, Grosse Frie K, Black N, van der Meulen J: Factors associated with non-response in routine use of patient reported outcome measures after elective surgery in England. *Health Qual Life Outcomes* 2012; 10: 34
- Judge A, Arden NK, Batra RN, Thomas G, Beard D, Javaid MK, et al.: The association of patient characteristics and surgical variables on symptoms of pain and function over 5 years following primary hip-replacement surgery: a prospective cohort study. *BMJ Open* 2013; 3: e002453
- Judge A, Cooper C, Williams S, Dreinhöfer K, Dieppe P: Patient-reported outcomes one year after primary hip replacement in a European Collaborative Cohort. *Arthritis Care Res* 2010; 62: 480–488
- Judge A, Cooper C, Arden NK et al.: Pre-operative expectation predicts 12-month post-operative outcome among patients undergoing primary total hip replacement in European orthopaedic centres. *Osteoarthritis Cartilage* 2011; 19: 659–67
- Judge A, Javaid MK, Arden NK et al.: Clinical tool to identify patients who are most likely to achieve long-term improvement in physical function after total hip arthroplasty. *Arthritis Care Res* 2012; 64: 881–9
- Judge A, Batra RN, Thomas GE et al.: Body mass index is not a clinically meaningful predictor of patient reported outcomes of primary hip replacement surgery: prospective cohort study. *Osteoarthritis Cartilage* 2014; 22: 431–9
- Jüni P, Low N, Reichenbach S, Villiger PM, Williams S, Dieppe PA: Gender inequity in the provision of care for hip disease: population-based cross-sectional study. *Osteoarthritis Cartilage* 2010; 18: 640–5
- Kapoor A, Chew P, Silliman RA et al.: Venous thromboembolism after joint replacement in older male veterans with comorbidity. *J Am Geriatr Soc* 2013; 61: 590–601
- Mancuso CA, Jout J, Salvati EA, Sculco TP: Fulfillment of patients' expectations for total hip arthroplasty. *The Journal of bone and joint surgery* 2009; American volume 91: 2073–78

31. Martinez-Huedo MA, Villanueva M, de Andres AL et al.: Trends 2001 to 2008 in incidence and immediate postoperative outcomes for major joint replacement among Spanish adults suffering diabetes. *Eur J Orthop Surg Traumatol* 2013; 23: 53–9
32. Neuburger J, Hutchings A, Black N, van der Meulen JH: Socioeconomic differences in patient-reported outcomes after a hip or knee replacement in the English National Health Service. *Journal of Public Health* 2013; 35: 115–24
33. Nilsson A: Predictors of patient relevant outcome after total hip replacement for osteoarthritis: a prospective study. *Annals of the Rheumatic Diseases* 2003; 62: 923–30
34. Palazzo C, Jourdan C, Descamps S et al.: Determinants of satisfaction 1 year after total hip arthroplasty: the role of expectations fulfilment. *BMC Musculoskeletal Disord* 2014; 15: 53
35. Peters ML, Sommer M, de Rijke JM et al.: Somatic and psychologic predictors of long-term unfavorable outcome after surgical intervention. *Ann Surg* 2007; 245: 487–94
36. Prokopetz JJ, Losina E, Bliss RL, Wright J, Baron JA, Katz JN: Risk factors for revision of primary total hip arthroplasty: a systematic review. *BMC Musculoskeletal Disord* 2012; 13: 251
37. Quintana JM, Escobar A, Aguirre U, Lafuente I, Arenaza JC: Predictors of Health-related Quality-of-life Change after Total Hip Arthroplasty. *Clin Orthop Relat Res* 2009; 467: 2886–94
38. Rolfson O, Dahlberg LE, Nilsson JA et al.: Variables determining outcome in total hip replacement surgery. *J Bone Joint Surg Br* 2009; 91: 157–61
39. Santaguida PL, Hawker GA, Hudak PL et al.: Patient characteristics affecting the prognosis of total hip and knee joint arthroplasty: a systematic review. *Can J Surg* 2008; 51: 428–36
40. Schäfer T, Pritzkeleit R, Jeszenszky C et al.: Trends and geographical variation of primary hip and knee joint replacement in Germany. *Osteoarthritis and Cartilage* 2013; 21: 279–88
41. Schäfer T, Krummenauer F, Mettelsiefen J, Kirschner S, Günther KP: Social, educational, and occupational predictors of total hip replacement outcome. *Osteoarthritis and Cartilage* 2010; 18: 1036–42
42. Schäfer T, Biakowski S, Walther A, Fickert S, Krummenauer F, Günther KP: Preoperative importance of improvement and postoperative satisfaction in patients undergoing total hip replacement. *Eur Orthop Traumatol* 2011; 1: 143–52
43. Singh JA, Lewallen DG: Ipsilateral lower extremity joint involvement increases the risk of poor pain and function outcomes after hip or knee arthroplasty. *BMC Med* 2013; 11: 144
44. Stiehler M, Goronzy J, Günther KP: Die endoprothetische Versorgung beim übergewichtigen Koxarthrosepatienten. *Orthopäde* 2015; 44: 523–30
45. Stürmer T, Dreinhöfer K, Gröber-Grätz D et al.: Differences in the views of orthopaedic surgeons and referring practitioners on the determinants of outcome after total hip replacement. *J Bone Joint Surg Br* 2005; 87: 1416–9
46. Tsang ST, Gaston P: Adverse peri-operative outcomes following elective total hip replacement in diabetes mellitus: a systematic review and meta-analysis of cohort studies. *Bone Joint J* 2013; 95-B: 1474–9
47. Vissers MM, Bussmann JB, Verhaar JA et al.: Psychological Factors Affecting the Outcome of Total Hip and Knee Arthroplasty: A Systematic Review. *Semin Arthritis Rheum* 2012; 41: 576–588
48. Young NL, Cheah D, Waddell JP, Wright JG: Patient characteristics that affect the outcome of total hip arthroplasty: a review. *Can J Surg* 1998; 41: 188–95
49. Zmistowski B1, Dizdarevic I, Jacovides CL, Radcliff KE, Mraovic B, Parvizi J: Patients with uncontrolled components of metabolic syndrome have increased risk of complications following total joint arthroplasty. *J Arthroplasty* 2013; 28: 904–7

# Vollständige Literatur zu Christian Dynybil: Rehabilitation nach Hüftendoprothese – was gibt es Neues? OUP 04-2017

## Literatur

- Hudak PL, Hogg-Johnson S, Bombardier C, McKeever PD, Wright JG: Testing a new theory of patient satisfaction with treatment outcome. *Med Care.* 2004; 42: 726–739
- Mahomed NN, Liang MH, Cook EF et al.: The importance of patient expectations in predicting functional outcomes after total joint arthroplasty. *J Rheumatol.* 2002; 29: 1273–1279
- Orbell S, Johnston M, Rowley D, Espley A, Davey P: Cognitive representations of illness and functional and affective adjustment following surgery for osteoarthritis. *Soc Sci Med.* 1998; 47: 93–102
- Kehlet H, Dahl JB: Anaesthesia, surgery, and challenges in postoperative recovery. *Lancet.* 2003; 362: 1921–1928
- Kehlet H, Wilmore D: Multimodal strategies to improve surgical outcome. *Am J Surg.* 2002; 183: 630–641
- Berger R: Podium Präsentation. Rush Medical Center Chicago. IL, AAOS 2016. Orlando/USA
- Dorr LD, Maheshwari AV, Long WT: Early pain relief and function after posterior minimally invasive and conventional total hip arthroplasty. A prospective, randomized, blinded study. *J Bone Joint Surg Am.* 2007; 89: 1153–1160
- Goebel S, Steinert AF, Schillinger J et al.: Reduced postoperative pain in total hip arthroplasty after minimal-invasive anterior approach. *Int Orthop.* 2012; 36: 491498
- Fink B, Mittelstaedt A, Schulz MS, Sebena P, Singer J: Comparison of a minimally invasive posterior approach and the standard posterior approach for total hip arthroplasty. A prospective and comparative study. *J Orthop Surg Res.* 2010; 5: 46
- Mouillhade F, Matsoukis J, Oger P, Mandereau C, Brzakala V, Dujardin F: Component positioning in primary total hip replacement: a prospective comparative study of two anterolateral approaches, minimally invasive versus gluteus medius hemimiotomy. *Orthop Traumatol Surg Res.* 2011; 97: 14–21
- Yang C, Zhu Q, Han Y et al.: Minimally-invasive total hip arthroplasty will improve early postoperative outcomes: a prospective, randomized, controlled trial. *Ir J Med Sci.* 2010; 179: 285–290
- Alecci V, Valente M, Crucil M, Minerva M, Pellegrino CM, Sabbadini DD: Comparison of primary total hip replacements performed with a direct anterior approach versus the standard lateral approach: perioperative findings. *J Orthop Traumatol.* 2011; 12: 123–129
- Jerosch J: Minimalinvasive Hüftendoprothetik. *Dt Ärzteblatt.* 2006; 103: 3333–3339
- Eingartner C: Current trends in total hip arthroplasty. *Ortho Traum Rehab.* 2007; 9: 8–14
- Oganda L, Wilson R, Archbold P et al.: A Minimal-Incision Technique in Total Hip Arthroplasty Does Not Improve Early Postoperative Outcomes. *J Bone Joint Surg Am.* 2005; 87: 701–710
- Spaans AJ, van den Hout JA, Bolder SB: High complication rate in the early experience of minimally invasive total hip arthroplasty by the direct anterior approach. *Acta Orthop.* 2012; 83: 342–346
- Smith TO, Blake V, Hing CB: Minimally invasive versus conventional exposure for total hip arthroplasty: a systematic review and meta-analysis of clinical and radiological outcomes. *Int Orthop.* 2011; 35: 173–184
- Lawlor M, Humphreys P, Morrow E, Ogonda L, Bennett D, Elliott D, Beverland D: Comparison of early postoperative functional levels following total hip replacement using minimally invasive versus standard incisions. A prospective randomized blinded trial. *Clin Rehabil.* 2005; 19: 465–474
- Goebel S, Steinert AF, Schillinger J et al.: Reduced postoperative pain in total hip arthroplasty after minimal-invasive anterior approach. *Int Orthop.* 2012; 36: 491–498
- Peck CN, Foster A, McLaughlan GJ: Reducing incision length or intensifying rehabilitation: what makes the difference to length of stay in total hip replacement in a UK setting? *Int Orthop.* 2006; 30: 395–398
- Pagnano MW, Trousdale RT, Meneghini RM, Hanssen AD: Slower recovery after two-incision than minimally-invasive total hip arthroplasty. A randomized clinical trial. *J Bone Joint Surg Am.* 2008; 90: 1000–1006
- Matta JM, Shahrdar C, Ferguson T: Single incision anterior approach for total hip arthroplasty on an orthopaedic table. *Clin Orthop Relat Res.* 2005; 441: 115–124
- Pagnano MW, Trousdale RT, Meneghini RM, Hanssen AD: Patients preferred a mini-posterior THA to a contralateral two-incision THA. *Clin Orthop Relat Res.* 2006; 453: 156–159
- Dorr LD, Thomas D, Long WT, Polatin PB, Sirianni LE: Psychologic reasons for patients preferring minimally invasive total hip arthroplasty. *Clin Orthop Relat Res.* 2007; 458: 94–100
- Malik A, Dorr LD: The science of minimally invasive total hip arthroplasty. *Clin Orthop Relat Res.* 2007; 463: 74–84
- Camu F, Beecher T, Recker DP, Verburg KM: Valdecoxib, a COX-2-specific inhibitor, is an efficacious, opioid-sparing analgesic in patients undergoing hip arthroplasty. *Am J Ther.* 2002; 9: 43–51
- Husted H, Hansen HC, Holm G et al.: What determines length of stay after total hip and knee arthroplasty? A nationwide study in Denmark. *Arch Orthop Trauma Surg.* 2010; 130: 263–268
- Sharma V, Morgan PM, Cheng EY: Factors influencing early rehabilitation after THA: a systematic review. *Clin Orthop Relat Res.* 2009; 467: 1400–1411
- Horlocker TT, Kopp SL, Pagnano MW, Hebl JR: Analgesia for total hip and knee arthroplasty: a multimodal pathway featuring peripheral nerve block. *J Am Acad Orthop Surg.* 2006; 14: 126–135
- Berger RA, Sanders SA, Thill ES, Sporer SM, Della Valle C: Newer anesthesia and rehabilitation protocols enable outpatient hip replacement in selected patients. *Clin Orthop Relat Res.* 2009; 467: 1424–1430
- Long WT, Ward SR, Dorr LD, Raya J, Boutary M, Sirianni LE: Postoperative pain management following total knee arthroplasty: a randomized comparison of continuous epidural versus femoral nerve infusion. *J Knee Surg.* 2006; 19: 137–143
- Maheshwari AV, Boutary M, Yun AG, Sirianni LE, Dorr LD: Multimodal analgesia without routine parenteral narcotics for total hip arthroplasty. *Clin Orthop Relat Res.* 2006; 453: 231–238
- Husted H, Holm G, Jacobsen S: Predictors of length of stay and patient satisfaction after hip and knee replacement surgery: fast-track experience in 712 patients. *Acta Orthop.* 2008; 79: 168–173
- Lunn TH, Husted H, Solgaard S et al.: Intraoperative local infiltration analgesia for early analgesia after total hip arthroplasty: a randomized, double-blind, placebo-controlled trial. *Reg Anesth Pain Med.* 2011; 36: 424–429



35. Parvataneni HK, Shah VP, Howard H, Cole N, Ranawat AS, Ranawat CS: Controlling pain after total hip and knee arthroplasty using a multimodal protocol with local periarticular injections: a prospective randomized study. *J Arthroplasty*. 2007; 22 (6 Suppl 2): 33–38
36. Andersen KV, Pfeiffer-Jensen M, Haraldsted V, Søballe K: Reduced hospital stay and narcotic consumption, and improved mobilization with local and intra-articular infiltration after hip arthroplasty: a randomized clinical trial of an intra-articular technique versus epidural infusion in 80 patients. *Acta Orthop*. 2007; 78: 180–186
37. Linke C, Jaschinski R, Heitmann T et al.: *Biomed* 2013; 1. Berliner Orthopädiereport
38. Mears DC, Mears SC, Chelly JE, Dai F, Vulakovich KL: THA with a minimally invasive technique, multi-modal anesthesia, and home rehabilitation: factors associated with early discharge? *Clin Orthop Relat Res*. 2009; 467: 1412–1417
39. Singelyn FJ, Ferrant T, Malisse MF, Joris D: Effect of intravenous patient-controlled analgesia with morphine, continuous epidural analgesia, and continuous femoral nerve sheath block on rehabilitation after unilateral total-hip arthroplasty. *Reg Anaesth Pain Med*. 2005; 30: 452–457
40. Becchi C, Al Malyan M, Coppini R, Campolo M, Magherini M, Boncinelli S: Opioid-free analgesia by continuous psoas compartment block after total hip arthroplasty. A randomized study. *Eur J Anaesthesiol*. 2008; 25: 418–423
41. Sharma V, Morgano PM, Cheng EY: Factors influencing early rehabilitation after THA: a systematic review. *Clin Orthop Relat Res*. 2009; 467: 1400–1411
42. Kehlet H, Slim K. The future of fast-track surgery. *Br J Surg*. 201; 99: 1025–1026
43. Ilfeld BM, Mariano ER, Girard PJ et al.: A multicenter, randomized, triple-masked, placebo-controlled trial of the effect of ambulatory continuous femoral nerve blocks on discharge-readiness following total knee arthroplasty in patients on general orthopaedic wards. *Pain*. 2010; 150: 477–484
44. McDonald S, Hetrick S, Green S: Pre-operative education for hip or knee replacement. *Cochrane Database Syst Rev*. 2004; (1): CD003526.
45. McDonald S, Page MJ, Behringer K, Wasik J, Sprowson A: Preoperative education for hip or knee replacement. *Cochrane Database Syst Rev*. 2014; (5): CD003526
46. Hozack W: Podium Präsentation, The Rothman Institute, Philadelphia, PA, AAOS 2016. Orlando/USA
47. Minns Lowe CJ, Barker KL, Dewey ME, Sackley CM: Effectiveness of physiotherapy exercise following hip arthroplasty for osteoarthritis: a systematic review of clinical trials. *BMC Musculoskelet Disord*. 2009; 10: 98
48. Minns Lowe CJ, Barker KL, Dewey M, Sackley CM: Effectiveness of physiotherapy exercise after knee arthroplasty for osteoarthritis: systematic review and meta-analysis of randomised controlled trials. *BMJ*. 2007; 335: 812
49. Husted H, Fast-track hip and knee arthroplasty: clinical and organizational aspects. *Acta Orthop Suppl*. 2012; 83: 1–39
50. Peter WF, Nelissen RG, Vlieland TP: Guideline recommendations for post-acute postoperative physiotherapy in total hip and knee arthroplasty: are they used in daily clinical practice? *Musculoskeletal Care*. 2014; 12: 125–131
51. Delorme TL: Restoration of muscle power by heavy-resistance exercises. *J Bone Joint Surg Am*. 1945; 27: 645–667
52. Delorme TL, Watkins AL. Technics of progressive resistance exercise. *Arch Phys Med Rehabil*. 1948 29: 263–273
53. Munin MC, Rudy TE, Glynn NW, Cressett LS, Rubash HE: Early inpatient rehabilitation after elective hip and knee arthroplasty. *JAMA*. 1998; 279: 847–852
54. Bandholm T, Kehlet H: Physiotherapy exercise after fast-track total hip and knee arthroplasty: time for reconsideration? *Arch Phys Med Rehabil*. 2012; 93: 1292–1294
55. Husby VS, Helgerud J, Bjørgen S, Husby OS, Benum P, Hoff J: Early maximal strength training is an efficient treatment for patients operated with total hip arthroplasty. *Arch Phys Med Rehabil*. 2009; 90: 1658–1667
56. Toigo M, Boutellier U: New fundamental resistance exercise determinants of molecular and cellular muscle adaptations. *Eur J Appl Physiol*. 2006; 97: 643–663
57. Okoro T, Lemmey AB, Maddison P, Andrew JG: An appraisal of rehabilitation regimes used for improving functional outcome after total hip replacement surgery. *Sports Med Arthrosc Rehabil Ther Technol*. 2012; 4–5
58. Lenssen AF, Crijs YH, Waltjé EM et al.: Efficiency of immediate postoperative inpatient physical therapy following total knee arthroplasty: an RCT. *BMC Musculoskelet Disord*. 2006; 7: 71
59. Moffet H, Collet JP, Shapiro SH, Paradis G, Marquis F, Roy L: Effectiveness of intensive rehabilitation on functional ability and quality of life after first total knee arthroplasty: A single-blind randomized controlled trial. *Arch Phys Med Rehabil*. 2004; 85: 546–556
60. Brower RG, Consequences of bed rest. *Crit Care Med*. 2009; 37(10 Suppl): 422–428
61. Piva SR, Teixeira PE, Almeida GJ et al.: Contribution of hip abductor strength to physical function in patients with total knee arthroplasty. *Phys Ther*. 2011; 91: 225–233
62. Toigo M, Boutellier U: New fundamental resistance exercise determinants of molecular and cellular muscle adaptations. *Eur J Appl Physiol*. 2006; 97: 643–663
63. Silkman Baker C, Mc Keon JM: Does pre-operative rehabilitation improve patient – based outcomes in persons who have undergone total knee arthroplasty? A systematic review. *PM R* 2012; 4: 756–767
64. Lenssen AF, de Bie RA: Role of physiotherapy in peri-operative management in total knee and hip surgery. *Injury*. 2006; 37 Suppl 5: 41–43
65. Khan F, Ng L, Gonzalez S, Hale T, Turner-Stokes L: Multidisciplinary rehabilitation programmes following joint replacement of the hip and knee in chronic arthropathy. *Cochrane Database Syst Rev*. 2008; (2): CD004957
66. Tayrose G, Newman D, Slover J, Jaffe F, Hunter T, Bosco J 3rd: Rapid mobilization decreases length-of-stay in joint replacement patients. *Bull Hosp Jt Dis*. 2013; 71: 222–226
67. Husby VS, Helgerud J, Bjørgen S, Husby OS, Benum P, Hoff J: Early postoperative maximal strength training improves work efficiency 6–12 months after osteoarthritis-induced total hip arthroplasty in patients younger than 60 years. *Am J Phys Med Rehabil* 2010; 89: 304–314
68. Kristensen J, Franklyn-Miller A. Resistance training in musculoskeletal rehabilitation: a systematic review. *Br J Sports Med*. 2012; 46: 719–726
69. Krank aus dem Krankenhaus. Visite. [www.ndr.de/Ratgeber/Gesundheit/Zufrueh-entlassen-krank-aus-demkrankenhaus](http://www.ndr.de/Ratgeber/Gesundheit/Zufrueh-entlassen-krank-aus-demkrankenhaus). Tina Roth. 26.05.2015
70. Greitemann B: REDIA-Studie. Die Welt, 2011
71. Heisel J, Jerosch J: Rehabilitationsfähigkeit – Rehabilitation nach Hüft- und Knieendoprothese. Köln: Deutscher Ärzte-Verlag, 2007: 71–73
72. Linke C, Jaschinski T, Heitmann T et al.: 1. Berliner Orthopädiereport. *Biomet*. 2013
73. Diehl R, Gebauer E, Groner A: *Kursbuch Sozialmedizin. Lehrbuch zum Curriculum der Bundesärztekammer*. Köln: Deutscher Ärzte-Verlag, 2012
74. Eiff W, Schüring S, Niehus C: REDIA. Auswirkungen der DRG – Einführung Rehabilitation. Berlin: Lit-Verlag, 2011
75. Statistisches Bundesamt. Grunddaten der Vorsorge- oder Rehabilitationseinrichtungen. Wiesbaden, 2009
76. Eiff W: Krankenhaus-IT Journal. Forschung. Ausgabe 03/2011
77. Reha-Bericht 2015, 6. Auflage, Deutsche Rentenversicherung Bund
78. Steiner M, Zwinkmann C, Riedel W et al.: Die medizinische Rehabilitation Erwerbstätiger – Sicherung von Produktivität und Wachstum. *Prognos AG* 2009
79. Eiff W, Schüring S: Kürzere Akutverweildauern erhöhen Aufwand in der Reha. *Dt. Ärzteblatt* 2011; 108: 1164–1166

# Vollständige Literatur zu Christian von Rüden et al.: Die Rolle des Weichteilschadens bei der Frakturbehandlung OUP 05-2017

## Literatur

- Tu YK, On Tong G, Wu CH, Sananpanich K, Kakinoki R: Soft-tissue injury in orthopaedic trauma. *Injury* 2008; 39 Suppl 4: 3–17
- Jordan DJ, Malahias M, Khan W, Hindocha S: The ortho-plastic approach to soft tissue management in trauma. *Open Orthop J* 2014; 8: 399–408
- Rittmann WW, Schibli M, Matter P, Allgower M: Open fractures. Long-term results in 200 consecutive cases. *Clin Orthop Relat Res* 1979; 138: 132–140
- Harry LE, Sandison A, Pearse MF, Paleolog EM, Nanchahal J: Comparison of the vascularity of fasciocutaneous tissue and muscle for coverage of open tibial fractures. *Plast Reconstr Surg* 2009; 124: 1211–1219
- Jaeger M, Maier D, Kern WV, Südkamp NP: Antibiotics in trauma and orthopedic surgery – a primer of evidence-based recommendations. *Injury* 2006; 37 Suppl 2: S74–80
- Court-Brown CM, Rimmer S, Prakash U, McQueen MM: The epidemiology of open long bone fractures. *Injury* 1998; 29: 529–534
- Gustilo RB, Mendoza RM, Williams DN: Problems in management of type III (severe) open fractures a new classification of type III open fractures. *J Trauma* 1984; 24: 742–746
- Tscherne H, Oestern HJ: Die Klassifizierung des Weichteilschadens bei offenen und geschlossenen Frakturen. *Unfallheilkunde* 1982; 85: 111–115
- Gregory RT, Gould RJ, Pecllet M, Wagner JS et al.: The mangled extremity syndrome: a severity grading system for multisystem injury of the extremity. *J Trauma* 1985; 25: 1147–1150
- Johansen K, Daines M, Howey T, Helfet D, Hansen ST Jr: Objective criteria accurately predicts amputation following lower extremity trauma. *J Trauma* 1990; 30: 568–573
- Howe HR Jr, Poole GV Jr, Hansen KJ et al.: Salvage of lower extremities following combined orthopedic and vascular trauma. A predictive salvage index. *Am Surg* 1987; 53: 205–208
- Russell WL, Sailors DM, Whittle TB, Fisher DF Jr, Burns RP: Limb salvage versus traumatic amputation. A decision based on a seven-part predictive index. *Ann Surg* 1991; 213: 473–481
- McNamara MG, Heckman JD, Corley FG: Severe open fractures of the lower extremity. A retrospective evaluation of the Mangled Extremity Severity Score (MESS). *J Orthop Trauma* 1994; 8: 81–87
- Ruedi TP, Buckley RE, Moran CG: *AO Principles of Fracture Management*. Vol 1-Principles 2nd Exp Ed. AO Publishing 2007: 91–111
- Byrd HS, Spicer TE, Cierny G: 3rd Management of open tibial fractures. *Plast Reconstr Surg* 1985; 76(5): 719–728
- Bosse MJ, MacKenzie EJ, Kellam JF et al.: A prospective evaluation of the clinical utility of the lower-extremity injury-severity scores. *J Bone Joint Surg Am* 2001; 83: 3–14
- Gustilo RB, Anderson JT: Prevention of infection in the treatment of one thousand and twenty-five open fractures of long bones retrospective and prospective analyses. *J Bone Joint Surg Am* 1976; 58: 453–458
- Gustilo RB, Gruninger RP, Davis T: Classification of type III (severe) open fractures relative to treatment and results. *Orthopedics* 1987; 10: 1781–1788
- Esterhai JL Jr, Queenan J: Management of soft tissue wounds associated with type III open fractures. *Orthop Clin North Am* 1991; 22: 427–432
- Gustilo RB, Merkow RL, Templeman D: The management of open fractures. *J Bone Joint Surg Am* 1990; 72: 299–304
- Standards for the management of open fractures of the lower limb, a short guide. *Guidelines Ed 1, Sept 2009*, British Orthopedic Association, British Association of Plastic Reconstructive and Aesthetic Surgeons
- Advanced Trauma Life Support (ATLS) for Doctors (2012). American College of Surgeons Committee on Trauma. 9th ed. Chicago/IL
- Kragh JF, O'Neill ML, Beebe DF et al.: Survey of the indications for use of emergency tourniquets. *J Spec Oper Med* 2011; 11: 30–38
- Rogers DM, Blouin GS, O'Leary JP: Povidone-iodine wound irrigation and wound sepsis. *Surg Gynecol Obstet* 1983; 157: 426–430
- Covey DC: Blast and fragment injuries of the musculoskeletal system. *J Bone Joint Surg Am* 2002; 84: 1221–1234
- Lerner A, Reis D, Soudry M: *Severe injuries to the limbs*. Staged treatment. Berlin, Heidelberg, New York: Springer; 2007, 233
- Yang EC, Eislner J: Treatment of isolated type I open fractures: is emergent operative debridement necessary. *Clin Orthop Relat Res* 2003; 410: 289–294
- Zalavras CG, Patzakis MJ: Open fractures evaluation and management. *J Am Acad Orthop Surg* 2003; 11: 212–219
- Roberts CS, Pape HC, Jones AL, Malkani AL, Rodriguez JL, Giannoudis PV: Damage control orthopaedics: evolving concepts in the treatment of patients who have sustained orthopaedic trauma. *Instr Course Lect* 2005; 54: 447–462
- Tuttle MS, Smith WR, Williams AE et al.: Safety and efficacy of damage control external fixation versus early definitive stabilization for femoral shaft fractures in the multiple-injured patient. *J Trauma* 2009; 67: 602–605
- Taeger G, Ruchholtz S, Waydhas C, Lewan U, Schmidt B, Nast-Kolb D: Damage control orthopedics in patients with multiple injuries is effective, time saving, and safe. *J Trauma* 2005; 59: 409–416
- Harwood PJ, Giannoudis PV, Probst C, Krettek C, Pape HC: The risk of local infective complications after damage control procedures for femoral shaft fracture. *J Orthop Trauma* 2006; 20: 181–189
- Kindsfater K, Jonassen EA: Osteomyelitis in grade II and grade III open tibia fractures with late debridement. *J Orthop Trauma* 1995; 9: 121–127
- Naique SB, Pearse M, Nanchahal J: Management of severe open tibial fractures the need for combined orthopaedic and plastic surgical treatment in specialist centres. *J Bone Joint Surg Br* 2006; 88: 351–357
- Singh J, Rambani R, Hashim Z, Raman R, Sharma HK: The relationship between time to surgical debridement and incidence of infection in grade III open fractures. *Strategies Trauma Limb Reconstr* 2012; 7: 33–37
- Pollak AN, Jones AL, Castillo RC, Bosse MJ, MacKenzie EJ, LEAP Study Group: The relationship between time to surgical debridement and incidence of infection after open high-energy lower extremity trauma. *J Bone Joint Surg Am* 2010; 92: 7–15
- Webb LX, Bosse MJ, Castillo RC, MacKenzie EJ, LEAP Study Group: Analysis of surgeon-controlled variables in the treatment of limb-threatening type-III open tibial diaphyseal fractures. *J Bone Joint Surg Am* 2007; 89: 923–928
- Harley BJ, Beaupre LA, Jones CA, Dulai SK, Weber DW: The effect of time to definitive treatment on the rate of non-

- union and infection in open fractures. *J Orthop Trauma* 2002; 16: 484–490
39. Ashford RU, Mehta JA, Cripps R: Delayed presentation is no barrier to satisfactory outcome in the management of open tibial fractures. *Injury* 2004; 35: 411–416
40. Skaggs DL, Friend L, Alman BM et al.: The effect of surgical delay on acute infection following 554 open fractures in children. *J Bone Joint Surg Am* 2005; 87: 8–12
41. Schenker ML, Yannascoli S, Baldwin KD, Ahn J, Mehta S: Does timing to operative debridement affect infectious complications in open long-bone fractures? A systematic review. *J Bone Joint Surg Am* 2012; 94: 1057–1064
42. Pfeifer R, Pape HC: Diagnostik und Versorgungsstrategien beim polytraumatisierten Patienten. *Chirurg* 2016; 87(2): 165–173
43. Lee JJ, Patel R, Biermann JS, Dougherty PJ: The musculoskeletal effects of cigarette smoking. *J Bone Joint Surg Am* 2013, 95: 850–859
44. White CB, Turner NS, Lee GC, Haidukewych GJ: Open ankle fractures in patients with diabetes mellitus. *Clin Orthop Relat Res* 2003; 414: 37–44
45. Gabel G, Pyrc J, Hinterseher I, Zwipp H, Saeger HD, Bergert H: Arterial injuries combined with open fractures – management and therapy. *Zentralbl Chir* 2009; 134: 292–297
46. Schlickewei W, Kuner EH, Mullaji AB, Gotze B: Upper and lower limb fractures with concomitant arterial injury. *J Bone Joint Surg Br* 1992; 74(2): 181–188
47. Gumbel D, Naundorf M, Napp M, Ekkernkamp A, Seifert J: Diagnostik und Management peripherer Gefäßverletzungen. *Unfallchirurg* 2014; 117: 445–459
48. Pape HC, Giannoudis P, Krettek C: The timing of fracture treatment in polytrauma patients: relevance of damage control orthopedic surgery. *Am J Surg* 2002; 183: 622–629
49. Blick SS, Brumback RJ, Poka A, Burgess AR, Ebraheim NA: Compartment syndrome in open tibial fractures. *J Bone Joint Surg Am* 1986; 68: 1348–53
50. World Society for Abdominal Compartment Syndrome Consensus: Definitions and Recommendations. <http://www.wsacs.org>
51. Weitz-Marshall AD, Bosse MJ: Timing of closure of open fractures. *J Am Acad Orthop Surg* 2002; 10: 379–384
52. Labler L, Trentz O: The use of vacuum assisted closure (VAC) in soft tissue injuries after high energy pelvic trauma. *Langenbecks Arch Surg* 2007; 392: 601–609
53. Labler L, Mica L, Härter L, Trentz O, Keel M: Influence of V.A.C.-therapy on cytokines and growth factors in traumatic wounds. *Zentralbl Chir* 2006; 131 Suppl 1: S62–67
54. von Rüden C, Benninger E, Mayer D, Trentz O, Labler L: Bogota-VAC – A Newly Modified Temporary Abdominal Closure Technique. *Eur J Trauma Emerg Surg* 2008; 34: 582–586
55. Patzakis MJ, Harvey JP Jr, Ivler D: The role of antibiotics in the management of open fractures. *J Bone Joint Surg Am* 1974; 56: 532–541
56. Lack WD, Karunakar MA, Angerame MR et al.: Type III Open Tibia Fractures: Immediate Antibiotic Prophylaxis Minimizes Infection. *J Orthop Trauma* 2015; 29: 1–6
57. Wacha H, Hoyme, U, Isenmann R et al.: Perioperative Antibiotika-Prophylaxe. Empfehlungen einer Expertenkommission der Paul-Ehrlich-Gesellschaft für Chemotherapie e.V. *Chemother J* 2010; 19: 70–84
58. Rhee P, Nunley MK, Demetriades D, Velmahos G, Doucet JJ: Tetanus and trauma: a review and recommendations. *J Trauma* 2005; 58: 1082–1088
59. [http://www.rki.de/DE/Content/Infekt/EpidBull/Merkblaetter/Ratgeber\\_Tetanus.html#doc2398266bodyText16](http://www.rki.de/DE/Content/Infekt/EpidBull/Merkblaetter/Ratgeber_Tetanus.html#doc2398266bodyText16)

# Vollständige Literatur zu Harald Hempfling: Die Patella partita, ein Apophysen-Äquivalent OUP 06-2017

## Literatur

1. Aams JD, Leonard RD: A developmental anomaly of the patella frequently diagnosed as fracture. *Surg Gynecol Obstet* 1925; 41: 601–604
2. Batten J, Menelaus MB: Fragmentation of the proximal pole of the patella. *J Bone Joint Surg Br* 1985; 67: 249–251
3. Baumgartl F: *Das Kniegelenk*. Berlin: Springer-Verlag, 1964
4. Bernhang AM, Levine SA: Familial absence of the patella. *J Bone Joint Surg Am* 1973; 55A:1088–1090
5. Blankstein A, Cohen I, Salai M, Diamant L, Chechick A, Ganel A: Ultrasonography: an imaging modality enabling the diagnosis of bipartite patella. *Knee Surg, Sports Traumatol, Arthrosc* 2001; 9: 221–224
6. Bohndorf K, Imhof H, Wörtler K: *Radiologische Diagnostik der Knochen und Gelenke (3. Auflage)*. Stuttgart: Thieme, 2014
7. Bourne MH, Bianco AJ: Bipartite patella in the adolescent: results of surgical excision. *J Pediatr Orthop* 1990; 10: 69–73
8. Caffey J: *Pediatric X-ray diagnosis*. Year Book Medical Publishers. Chicago, 1978: 1095
9. Canizares GH, Selesnick FH: Bipartite patella fracture. *Arthroscopy* 2003; 19: 215–217
10. Carneiro M, de Souza Nery CA, Mestriner LA: Bilateral stress fracture of the patellae: A case report. *The Knee* 2006; 13: 164–166
11. Carter SR: Traumatic separation of a bipartite patella. *Injury* 1989; 20: 244
12. Cipolla M, Cerulle G, Franco V, Gianni E, Puddo G: The double patella syndrome. *Knee Surg Sports Traumatol Arthroscopy* 1995; 3: 21–25
13. Devas MB: Stress fractures of the patella. *J Bone Joint Surg* 1960; 42B: 71
14. Douarre M: Anomalies d'ossification de la rotule. *Bull Soc Nat Chir* 1921; 8: 10
15. Dowd GSE: Marginal fractures of the patella. *Injury* 1982; 14: 287–291
16. Echeverria TS, Bersani FA: Acute fracture simulating a symptomatic bipartite patella. *Am J SportsMed* 1980; 8: 48–50
17. Ficat RP, Hungerford DS: *Disorders of the Patella-femoral Joint*. Baltimore: Williams and Wilkins, 1977
18. Finsterbush A, Lowe J, Haymsky G, Mann G: Symtomatic bipartite patella and related conditions. *Arthroscopy* 1999; 15/1, Suppl I, AGA-Abstracts
19. Finsterbush A, Lowe J, Haymsky G, Mann G: Symptomatic bipartite patella and related conditions. *Aus 5th Turkish Sports Traumatology Arthroscopy and Knee Wurgery Congress Istanbul* 24. to 27. Oct. 2000
20. Fründ H: Traumatische Chondropathie der Patella, ein selbständiges Krankheitsbild. *Zbl Chir* 1926; 53: 707
21. George R: Bilateral bipartite patellae. *Br J Surg* 1935; 22: 555–560
22. Goergen TG, Resnick D, Greenway G, Salzstein SL: Dorsal defect of the patella: A characteristic radiographic lesion. *Radiology* 1979; 130: 333
23. Green WT: Painful bipartite patellae. A report of three cases. *Clin Orthop* 1975; 110: 197–200
24. Gruber W: In Bildungsanomalie mir Bildungshemmung begründete Bipartition beider Patellae eines jungen Subjektes. *Virchows Arch Path Anat* 1983; 94: 358
25. Haenisch F: Verdoppelung der Patella in sagittaler Richtung. *Fortschr Röntgenstr* 1925; 33: 678
26. Haswell DM, Berne AS, Graham CB: The dorsal defect of the patella. *Pediatr Radiol* 1976; 4: 238
27. Halpern AA, Hewitt O: Painful medial bipartite patellae. *Clin Orthop Relat Res* 1978; 134: 180–181
28. Hems T, Tillmann B: Tendon entheses off he human masticatory muscles. *Anat Embryol* 2000; 202: 201–208
29. Iossifidis A, Brueton RN, Nunan TO: Bone-scintigraphy in painful bipartite patella. *Eur J Nucl Med* 1995; 22: 1212–1213
30. Iossifidis A, Brueton RN: Painful bipartite patella following injury. *Injury* 26 1995; 175–176
31. Ireland ML, Chang JL: Acute fracture bipartite patella: case report and literature review. *Med Sci Sports Exerc* 1995; 27: 299–302
32. Joachimsthal G: Über Struktur, Lage und Anomalien der menschlichen Kniescheibe. *Langenbecks Arch Klin Chir* 1902; 67: 342
33. Johansson S: Eine bisher nicht beschriebene Erkrankung der Patella. *Hygiea (Stockh)* 1922; 84: 161
34. Jones B: Congenital absence of both patellae. *J Bone Joint Surg Br* 1955; 37B: 352
35. Jünemann H, Schilling H: Schalenförmige Knochenapophyse an der Patella beidseits – eine seltene Formvariante des heranwachsenden Skelettes. *Akt Traumatol* 1984; 14: 101–102
36. Kavanagh EC, Zoga A, Omar I, Ford S, Schweitzer M, Eustace S: MRI findings in bipartite patella. *Skeletal Radiol* 2007; 36: 209–214
37. Kaya DO, Toprak U, Baltaci, Yosmapglu B, Ozer H: Long-term functional and sonographic outcomes in Osgood-Schlatter disease. *Knee Surg Sports Traumatol Arthrosc* 2013; 21: 1131–1139
38. Kempson FC: Emargination of the patella. *J Anat (Lond)* 1902; 36: 419
39. Lapidus PW: Longitudinal fractures of the patella. *J Bone Joint Surg* 1932; 14: 351
40. Lawson JP: Symptomatic radiographic variants in extremities. *Radiology* 1985; 157: 625
41. Linssen M, Hille E, Schulitz KP: Ist die Patella bipartita ein Ermüdungsgeschehen? *Orthop Praxis* 1988; 5: 294–295
42. Little EM: Congenital absence or delayed development of the patella. *Lancet* 1897; 2: 781–784
43. Mayer HN: Congenital absence or delayed development of the patella. *Lancet* 1897; 2: 1384–1385
44. Moreau L: Les fractures méconnues de la rotule. *Presse Méd* 1920; 28: 374
45. Ogata K: Painful bipartite patella. A new approach to operative treatment. *J Bone Joint Surg* 1994; 76: 573–578
46. Ogden JA, Mc Carthy SM, Jokl P: The painful bipartite patella. *J Pediatr Orthop* 1982; 2: 263
47. Ogden JA: Radiology of postnatal skeletal development. X. Patella and tibial tuberosity. *Skeletal Radiol* 1984; 11: 246–257
48. Ogden JA: *Skeletal injury in the child*. (3. Ed.). Heidelberg: Springer, 2000
49. Ogden JA: *Skeletal injury in the child*. (3. Ed.). Philadelphia: WB Saunders, 2000
50. Olbrich E: Patella emarginata – Patella partita. In: *Forschungen und Forscher der Tiroler Ärzteschule, II. Band (1948–1950)*
51. Oohashi Y, Noriki S, Koshino T, Fukuda M: Histopathological abnormalities in painful bipartite patellae in adolescents. *The Knee* 2006; 13: 189–193
52. Oohashi Y, Koshino T: Bone scintigraphy in patients with bipartite patella. *Knee Surg Sports Traumatol Arthrosc* 2007; 15: 1395–1399
53. Oohashi Y, Koshino T, Oohashi Y: Clinical features and classification of bip-

- artite or tripartite patella. *Knee Surg Sports Traumatol Arthrosc* 2010; 18: 1465–1469
54. Osgood RB: Lesions of the tibial tubercle occurring during adolescence. *Boston Med Surg J* 1903; 148: 114–117
55. Petty MJ: Two cases of abnormal Patellae. *British J Surg* 1928; 12: 799–800
56. Pfeil E: *Das Femoropatellargelenk*. Leipzig: Johann Ambrosius Barth, 1981
57. Prakash S, Chopra SRK, Indarji T: Traumatic separation of a bipartite patella. *Injury: The British Journal of Accident Surgery* 1989; 20: 244
58. Resnick D, Kang HS, Pretteklieber ML: *Internal Derangements of Joints* (2. Ed.). Philadelphia: Elsevier, 2007
59. Rohleder K: L'arthrose de la surface articulaire de la rotule. *Rev Chir Orthop* 1964; 50: 361
60. Rubin G: Congenital absence of patellae and other patellar anomalies in three members of the same family. *JAMA* 1915; 64: 2062
61. Saupe H: Beiträge zur Patella Bipartita. *Fortschr Geb Röntgenstr Nuklearmed Ergänzungsbad* 1921; 28: 37–46
62. Saupe E: Primäre Knochenmarkseiterung der Kniescheibe. *Deutsch Z Chir* 1943; 258: 386–392
63. Scapinelli R: Blood supply of the human patella. Ist relationto ischaemic necrosis after fracture. *J Bone Joint Surg* 1967; 49B: 563
64. Schaer H: Die Patella partita. *Ergebn Chir Orthop* 1934; 27: 1
65. Schlatterer C: Verletzungen des schnabelförmigen Fortsatzes der oberen Tibiaepiphyse. *Beitr Klin Chir* 1903; 38: 874–887
66. Scuderi GR: *The Patella*. Heidelberg: Springer, 1995
67. Simens W: Patella partita. *Dtsch Z Chir* 1931; 233: 727–755
68. Sillanpää PJ, Paakkala A, Paakkala T, Mäenpää H, Toivanen J: Displaced longitudinal stress fracture of the patella. *J Bone Joint Surg Am* 2010; 92: 2344–2347
69. Sinding-larsen C: A hitherto unknown affection of the patella in children. *Acta Radiol (Stockh)* 1921; 1: 171
70. Sonnenschein A: *Biologie, Pathologie und Therapie der Gelenke*. Dargestellt am Kniegelenk. Basel: Benno Schwabe & Co., 1952
71. Stocker RL, van Laer L: Injury of a bipartite patella in a young upcoming sportsman. *Arch Orthop Trauma Surg* 2011; 131: 75–78
72. Stoller DW: *Magnetic Resonance Imaging in Orthopaedics and Sports Medicine*. Philadelphia: Lippincott Williams & Wilkins, 2007
73. Tauber M, Matis N, Resch H: Traumatic separation of an uncommon bipartite patella type: a case report. *Knee Surg Sports Traumatol Arthrosc* 2007; 15: 83–87
74. Tillmann B, Peters A: Belastung und Beanspruchung der Patella. In: Hasselpflug J: *Die Blauth-Knieendoprothese: Grundlagen, gegenwärtiger Stand und Ausblick*. Bern: Huber, 1992: 54–59
75. Todd T, Mc Cally WC: Defects of the patellar border. *Ann Surg* 1921; 74: 775–782
76. Tos L, Salvi V: *La patolgia non traumatico della rotula*. Torino: Es Minerva Med 1968; 79–98
77. Trillat A: *Chirurgie du genu*. Lyon: Simep Editeur, 1976
78. Vahlensieck M, Reiser M: *MRT des Bewegungsapparates*. Stuttgart: Thieme, 2015
79. Van Hoenacker FM, Bernaerts A, Van de Perre S, de Scherper AM: Images in clinical radiology. *JBR-BRT* 2002; 85: 219
80. Van Holsbeeck, Vandamme B, Mkar-chal G, Martens M, Victor J, Baert AL: Dorsal defect of the patella: concept of its origin and relationship with bipartite and multipartite patella. *Skeletal Radiol* 1987; 16: 304–311
81. Weaver JK: Bipartite patellae as a cause of disability in the athlete. *Am J Sports Med* 1977; 5: 137–143
82. Ytrehus B, Haga HA, Mellum CN et al.: Experimental ischemia of porcine growth cartilage produces lesions of osteochondrosis. *J Orthop Res* 2004; 22: 1201–1209
83. Ytrehus B, Carlson CS, Ekma S: Etiology and pathogenesis of osteochondrosis. *Vet Pathol* 2007; 44: 429–448
84. Zumstein M, Sukthankar A, Exner GU: Tripartite patella: late appearance of a third ossification center in childhood. *J Pediatr Orthop B* 2006; 15: 75–76

# Vollständige Literatur zu Armin Keshmiri et al.: Die Patella in der Endoprothetik OUP 06-2017

## Literatur

1. Akagi M, Oh M, Nonaka T et al.: An Anteroposterior Axis of the Tibia for Total Knee Arthroplasty. *Clin. Orthop* 2004; 420
2. Bellemans J, Colyn W, Vandenuecker H, Victor J: The Chitranjan Ranawat award: is neutral mechanical alignment normal for all patients? The concept of constitutional varus. *Clin Orthop Relat Res.* 2012; 470: 45–53
3. Berger RA, Crosse LS, Jacobs JJ, Rubash HE: Malrotation Causing Patellofemoral Complications After Total Knee Arthroplasty. *Clin. Orthop. Relat. Res.* 1998; 356: 144–53
4. Callaghan MJ, Oldham JA, Winstanley J: A comparison of two types of electrical stimulation of the quadriceps in the treatment of patellofemoral pain syndrome: A pilot study. *Clin Rehabil* 2001; 15: 637–46
5. Cerciello S, Robin J, Lustig S, Maccauro G, Heyse TJ, Neyret P: The role of patelloplasty in total knee arthroplasty. *Arch Orthop Trauma Surg.* 2016; 136: 1607–13
6. France L, Nester C: Effect of errors in the identification of anatomical landmarks on the accuracy of Q angle values. *Clinical biomechanics* (Bristol, Avon). 2001; 16: 710–3
7. Gromov K, Korchi M, Thomsen MG, Husted H, Troelsen A: What is the optimal alignment of the tibial and femoral components in knee arthroplasty? *Acta Orthop.* 2014; 85: 480–7
8. Heinert G, Kendoff D, Preiss S et al.: Patellofemoral kinematics in mobile-bearing and fixed-bearing posterior stabilised total knee replacements: a cadaveric study. *Knee Surg Sports Traumatol Arthrosc* 2011; 19: 967–972
9. Howell SM, Howell SJ, Kuznik KT, Cohen J, Hull ML: Does a kinematically aligned total knee arthroplasty restore function without failure regardless of alignment category? *Clin Orthop Relat Res.* 2013; 471: 1000–7
10. Johnson LL, van Dyk GE, Green JR 3rd, Pittsley AW, Bays B, Gully SM, Phillips JM: Clinical assessment of asymptomatic knees: comparison of men and women. *Arthroscopy.* 1998; 14: 347–59
11. Keshmiri A, Springorum HR, Baier C, Zeman F, Grifka J, Maderbacher G: Changes in sagittal component alignment alters patellar kinematics in TKA: an in vitro study. *Knee Surg Sports Traumatol Arthrosc.* 2016; 24: 823–9
12. Keshmiri A, Springorum H, Baier C, Zeman F, Grifka J, Maderbacher G: Is it possible to re-establish pre-operative patellar kinematics using a ligament-balanced technique in total knee arthroplasty? A cadaveric investigation. *Int Orthop.* 2015; 39: 441–8
13. Keshmiri A, Maderbacher G, Baier C, Müller W, Grifka J, Springorum HR: Do surgical patellar interventions restore patellar kinematics in fixed-bearing, cruciate-retaining total knee arthroplasty? An in vitro study. *J Arthroplasty.* 2014; 29: 2197–201
14. Kienapfel H, Springorum HP, Ziegler A et al.: Der Einfluss der Femur- und Tibiakomponentenrotation auf das patellofemorale Versagen beim künstlichen Kniegelenkersatz. *Orthop* 2003; 32: 312–318
15. Kolowich PA, Paulos LE, Rosenberg TD, Farnsworth S: Lateral release of the patella: indications and contraindications *Am J Sports Med.* 1990; 18: 359–65
16. Li B, Bai L, Fu Y, Wang G, He M, Wang J: Comparison of clinical outcomes between patellar resurfacing and nonresurfacing in total knee arthroplasty: retrospective study of 130 cases. *J Int Med Res.* 2012; 40: 1794–803
17. Luyckx T, Didden K, Vandenuecker H, Labey L, Innocenti B, Bellemans J: Is there a biomechanical explanation for anterior knee pain in patients with patella alta?: influence of patellar height on patellofemoral contact force, contact area and contact pressure. *J Bone Joint Surg Br.* 2009; 91: 344–50
18. Maderbacher G, Baier C, Benditz A, Wagner F, Greimel F, Grifka J, Keshmiri A: Presence of rotational errors in long leg radiographs after total knee arthroplasty and impact on measured lower limb and component alignment. *Int Orthop.* 2017; Jan 31. doi: 10.1007/s00264-017-3408-3. (Epub ahead of print)
19. Maderbacher G, Keshmiri A, Schaumburger Jet al.: What is the optimal valgus pre-set for intramedullary femoral alignment rods in total knee arthroplasty? *Knee Surg Sports Traumatol Arthrosc* 2016; May 6 (epub ahead of print)
20. Maderbacher G, Keshmiri A, Springorum HR, Maderbacher H, Grifka J, Baier C: Influence of Component Rotation in Total Knee Arthroplasty on Tibio-Femoral Kinematics – a cadaveric investigation. *J Arthroplasty* 2017; DOI: <http://dx.doi.org/10.1016/j.arth.2017.03.055> (in press)
21. Maderbacher G, Schaumburger J, Baier C et al.: Appropriate sagittal femoral component alignment cannot be ensured by intramedullary alignment rods. *Knee Surg Sports Traumatol Arthrosc.* 2016; 24: 2453–60
22. Maderbacher G, Keshmiri A, Springorum HR, Greimel F, Grifka J, Baier C: Are We Subluxating Knees in Total Knee Arthroplasty? A Cadaveric Investigation. *J Arthroplasty* 2017; 32: 263–269
23. Miller MC, Berger RA, Petrella AJ et al.: Optimizing Femoral Component Rotation in Total Knee Arthroplasty. *Clin. Orthop.* 2001; 392
24. Pagenstert G, Seelhoff J, Henninger HB, Wirtz DC, Valderrabano V, Barg A: Lateral patellar facetectomy and medial reefing in patients with lateral facet syndrome after patellar-retaining total knee arthroplasty. *J Arthroplasty.* 2014; 29: 2156–62
25. Post W, Dye S: Patellofemoral pain: an enigma explained by homeostasis and common sense. *American Journal of Orthopedics* 2017; 46: 92–100
26. Singerman R, Davy DT, Goldberg VM: Effects of patella alta and patella infera on patellofemoral contact forces. *J Biomech.* 1994; 27: 1059–65
27. Staheli LT, Corbett M, Wyss C, King H: Lower-Extremity rotational problems in children. Normal values to guide management. *J Bone Joint Surg Am* 1984; 67: 39–47
28. Viktor J: Rotational alignment of the distal femur: a literature review. *Orthop Traumatol Surg Res.* 2009; 95: 365–72
29. Ward SR, Terk MR, Powers CM: Patella alta: association with patellofemoral alignment and changes in contact area during weight-bearing. *J Bone Joint Surg Am.* 2007; 89: 1749–55

# Vollständige Literatur zu Daniel Wagner: Patellofemorales Schmerzsyndrom OUP 06-2017

## Literatur

- Ahmed AM, Burke DL, Hyder A: Force analysis of the patellar mechanism. *J Orthop Res.* 1987; 5: 69–85
- Becher C, Schumacher T, Fleischer B, Ettinger M, Smith T, Ostermeier S: The effects of a dynamic patellar realignment brace on disease determinants of patellofemoral instability in the upright weight-bearing condition. *J Ortho Surg Res* 2015; 10: 126
- Boling MC, Bolgla LA, Mattacola CG, Uhl TL, Hosey RG: Outcomes of a weight-bearing rehabilitation program for patients diagnosed with patellofemoral pain syndrome. *Arch Phys Med Rehabil.* 2006; 87: 1428–35
- Brogden RN, Heel RC, Pakes GE, Spreight TM, Avery GS: Diflunisal: a review of its pharmacological properties and therapeutic use in pain and musculoskeletal strains and sprains and pain in osteoarthritis. *Drugs* 1980; 19: 84–106
- Brosseau L, Casimiro L, Robinson V et al.: Therapeutic ultrasound for treating patellofemoral pain syndrome: Intervention Review. *The Cochrane Library* 2001; (4): CD003375
- Callaghan MJ, Selve J, Bagley PJ, Oldham JA.: The effects of patellar taping on knee joint proprioception. *J Athl Train.* 2002; 37(1): 19–24
- Callaghan MJ, Oldham JA, Winstanley J: A comparison of two types of electrical stimulation of the quadriceps in the treatment of patellofemoral pain syndrome: A pilot study. *Clin Rehabil* 2001; 15: 637–46
- Can F et al.: Rehabilitation of patellofemoral pain syndrome: TENS versus diadynamic current therapy for pain relief. *The Pain Clinic.* 2003; 15: 61–8
- Christou EA: Patellar taping increases vastus medialis oblique activity in the presence of patellofemoral pain. *J Electromyogr Kinesiol* 2004; 14: 495–504
- Collins N, Crossley K, Beller E, Darnell R, McPoil T, Vicenzino B: Foot Orthoses and physiotherapy in the treatment of patellofemoral pain syndrome: randomised clinical trial. *Br J Sports Med;* 2009; 43: 169–71
- Conchie H, Clark D, Metcalfe A, Eldridge J, Whitehouse M: Adolescent knee pain and patellar dislocation are associated with patellofemoral osteoarthritis in adulthood: a case control study. *Knee* 2016; 23: 708–11
- Cowan SM, Bennell KL, Hodges PW: Therapeutic patellar taping changes the timing of vasti muscle activation in people with patellofemoral pain syndrome. *Clin J Sports Med;* 2002; 12: 339–47
- Cowan SM, Crossley KM, Bennell KL: Altered hip and trunk muscle function in individuals with patellofemoral pain. *Br J Sports Med* 2008; 43: 584–88
- Crossley KM, Marino GP, Macilquham MD, Schache AG, Hinman RS: Can patellar tape reduce the patellar malalignment and pain associated with patellofemoral osteoarthritis? *Arthritis Rheum* 2009; 61: 1719–25
- Darracott J: Treatment of the painful knee fulfilling diagnostic criteria for „chondromalacia patellae“. *Med Res&Opin* 1973; 1: 412–22
- De Anrade J et al.: Joint distension and reflex inhibition in the knee. *J Bone Joint Surg* 1965; 47a: 3131–22
- Dejour H, Walch G, Nove-Josserand L, Guier C: Factors of patellar instability: an anatomic radiographic study. *Knee Surg Sports Traumatol Arthrosc* 1994; 2: 19–26
- Devitt M: Acupuncture relieves anterior knee pain. *Acup Today* 2000; 1: 4
- D'hondt NE, Struijs PA, Kerkhoffs GM et al.: Orthotic devices for treating patellofemoral pain syndrome. *Cochrane Database Syst Rev* 2002; (2): CD002267
- Drake DF, Pidoce PE, Sricksen J: Botulinum toxin A for nonsurgical lateral release in patellofemoral pain syndrome: A case study. *Military Medicine* 2011; 6: 696–98
- Draper CE, Besier TF, Santos JM, Jennings F, Fredericson M, Gold GE et al.: Using real-time MRI to quantify altered joint kinematics in subjects with patellofemoral pain and to evaluate the effects of a patellar brace or sleeve on joint motion. *J Orthop Res.* 2009; 27: 571–7
- Dubin DCJ: Evidence based treatment for iliotibial band friction syndrome: Review of literature. *Sports Therapy* 2006; December
- Dye SF: The pathophysiology of patellofemoral pain: a tissue homeostasis perspective. *Clin Orthop Relat Res.* 2005; 436: 100–10
- Eckstein F, Lemberger B, Gratzke C et al.: In vivo cartilage deformation after different types of activity and its dependence on physical training status. *Ann Rheum Dis* 2005; 64: 291–95
- Eng JJ, Pierrynowski MR: Evaluation of soft foot orthotics in the treatment of patellofemoral pain syndrome. *Phys Ther.* 1993; 73: 62–8
- Fuentes JP et al.: Effectiveness of interferential current therapy in the management of musculoskeletal pain: a systematic review and metaanalysis. *Phys Ther* 2010; 90: 1219–38
- Fulkerson JP, Folcik MA: Comparison of diflunisal and naproxen for pain relief of anterior knee pain. *Clin Ther* 1986; 9: 59–61
- Gilleard W, McConnell J, Parsons D: The effect of patellar taping on the onset of vastus medialis obliquus and vastus lateralis muscle activity in persons with patellofemoral pain. *Phys Ther.* 1998; 78: 25–32
- Fredericson M, Powers CM: Practical management of patellofemoral pain. *Clin J Sports Med* 2002; 12: 36–38
- Heintjes EM et al.: Pharmacotherapy for patellofemoral pain syndrome (Review). *The Cochrane Library* 2008
- Herrington L, Pearson S: Does level of load affect relative activation levels of vastus medialis oblique and vastus lateralis? *J Electromyogr Kinesiol* 2005; 16: 379–83
- Herrington L et al.: The effect of limb position, exercise mode and contraction type on overall activity of VMO and VL. *Phys Ther Sport* 2006; 7: 87–92
- Jensen R, Goethesen O, Listh K, Baerheim A: Acupuncture treatment of patellofemoral pain syndrome. *J Altern Compl Med* 1999; 5: 521–27
- Kannus P, Natri A, Niitymäki S, Järvinen M: Effects of intraarticular glucosaminoglycan polysulfate treatment on patellofemoral pain syndrome. A prospective, randomized double-blind trial comparing GAG with placebo and quadriceps muscle exercises. *Arthritis Rheum* 1992; 35: 1035–61
- Lake D et al.: Effect of therapeutic modalities on patients with patellofemoral pain syndrome: a systematic review. *Sports Health* 2011; 3: 182–89
- Lowry CD, Cleland JA, Dyke K: Management of patients with patellofemoral pain syndrome using a multimodal approach: A case series. *J Orthop Sports Phys Ther* 2008; 38: 691–702
- Merican AM, Amis AA: Iliotibial band tension affects patellofemoral and tibiofemoral kinematics. *J Biomech.* 2009; 42: 1539–46

38. Nyland J et al.: Femoral anteversion influences vastus medialis and gluteus medius EMG amplitude: composite hip abductor EMG amplitude ratios during isometric combined hip abduction-external rotation. *J Electromyogr Kinesiol* 2003; 14: 255–61c
39. Pagenstert GI, Bachmann M. Klinische Untersuchung bei patellofemoralem Problem. *Orthopädie*. 2008; 37: 890–903
40. Pal S, Draper CE, Fredericson M, Gold GE, Delp SL, Beaupre GS, Besier TF: Patellar maltracking correlates with vastus medialis activation delay in patellofemoral pain patients. *Am J Sports Med*. 2011; 39: 590–8
41. Powers CM, Ward SR, Chan LD, Chen YJ, Terk MR: The effect of bracing on patella alignment and patellofemoral joint contact areas. *Med Sci Sports Exerc*; 2004; 36: 1226–32
42. Puniello MS: iliotibial band tightness and medial patellar glide in patients with patellofemoral dysfunction. *J Orthop Sports Phys Ther*. 1993; 17: 144–148
43. Raatikainen T et al.: Effect of glucosaminoglycan polysulfate on chondromalacia patellae: a placebo controlled 1 year study. *Acta Orth Scand* 1990; 42: 223–8
44. Reilly DT, Martens M: Experimental Analysis of the Quadriceps Muscle Force and Patello-Femoral Joint Reaction Force for Various Activities. *Acta Orthop Scand* 1972; 43: 126–37
45. Rowlands BW, Brantingham JW: The efficacy of patella mobilization in patients suffering from patellofemoral pain syndrome. *J Neuromusc Sys* 1999; 7: 142–49
46. Saleh KJ, Arendt EA, Eldridge J, Fulkeron JP, Minas T, Mulhall KJ: Symposium. Operative treatment of patellofemoral arthritis. *J Bone Joint Surg Am* 2005; 87: 859–67
47. Santos TRT, Oliveira BA, Ocarino JM, Holt KG, Fonseca ST: Effectiveness of hip muscle strengthening in patellofemoral pain syndrome patients: a systematic review. *Braz. J. Phys. Ther*. 2015; 19
48. Seitlinger G, Scheurecker G, Högler R, Kramer J, Hofmann S. Bildgebende Diagnostik des Patellofemoralgelenks. *Arthroskopie* 2010; 23: 176–83
49. Spencer et al.: Knee joint effusion and quadriceps reflex inhibition in man. *Arch Phys Med Rehabil* 1984; 65: 171–5
50. Singer BJ, Silbert PL, Song S, Dunne JW, Singer KP: Treatment of refractory anterior knee pain using botulinum toxin type A (Dysport) injection to the distal vastus lateralis muscle: a randomised placebo controlled crossover trial. *Br J Sports Med*. 2011; 45: 640–5
51. Steinkamp LA, Dillingham MF, Markel MD, Hill JA, Kaufman KR. Biomechanical considerations in patellofemoral joint rehabilitation. *Am J Sports Med* 1993; 21: 438–44
52. Stradford P: Electromyography of the quadriceps femoris muscles in subjects with normal knees and acutely effused knees. *Physical Ther* 1981; 62: 279–83
53. Suter E et al.: Conservative lower back pain treatment reduces inhibition in knee extensor muscle: a randomized controlled trial. *Manipulativ Physiol Ther* 2000; 23: 76–80
54. Thomas MJ, Wood L, Selfe J, Peat G: Anterior knee pain in younger adults as a precursor to subsequent patellofemoral osteoarthritis: a systematic review. *MBC Muskuloskelet Disord* 2010; 11: 201–8
55. Tyler TF, Nicholas SJ, Mullaney MJ, McHugh MP: The role of hip muscle function in the treatment of patellofemoral pain syndrome. *Am J Sports Med*. 2005; 34: 630–6
56. Utting MR, Davies G, Newman JH: is anterior knee pain a predisposing factor to patellofemoral osteoarthritis? *Knee* 2005; 12: 362–365
57. Van Mechelen W, Hlobil H, Kemper HCG: Incidence, severity, aetiology and prevention of sports injuries. *Sports Med* 1992; 14: 82–99
58. Vane JR. Inhibition of prostaglandin synthesis as a mechanism of action for aspirin-like drugs. *Nat New Biol*. 1971; 231: 232–5
59. Wilse HH, Fiebert I, Kates JL: EMG Biofeedback as treatment for patellofemoral pain syndrome. *J Orthop Sports Phys Ther* 1984; 6: 95–103
60. Wilson NA, Press JM, Koh JL, Hendrix RW, Zhang LQ: In vivo noninvasive evaluation of abnormal patellar tracking during squatting in patients with patellofemoral pain. *J Bone Joint Surg Am* 2009; 91: 558–566
61. Wilson NA, Mazahery BT, Koh JL, Zhang LQ: Effect of bracing on dynamic patellofemoral contact mechanics. *J Rehabil Res Dev*. 2010; 47: 531–41
62. Wu CC, Chou SW, Hong WH: Effectiveness of goosetep training or its modification on treating patellar malalignment syndrome: clinical, radiographic, and electromyographic studies. *Arch Orthop Trauma Surg*. 2009; 129: 333–41